

**A PROSPECTIVE STUDY OF CIGARETTE SMOKING AND RISK OF  
INCIDENT HYPERTENSION IN WOMEN**

Thomas S. Bowman, MD, MPH, J. Michael Gaziano, MD, MPH, Julie E. Buring ScD  
and Howard D. Sesso ScD, MPH

From the Veterans Affairs Boston Healthcare System - Massachusetts Veterans Affairs  
Epidemiology, Research, and Information Center (TSB, JMG); Divisions of Aging (TSB,  
JMG, JEB, HDS) and Preventive Medicine (JMG, JEB, HDS), Department of Medicine,  
Brigham and Women's Hospital; Department of Epidemiology (JEB, HDS), Harvard  
School of Public Health; and Department of Ambulatory Care and Prevention, Harvard  
Medical School (JEB) - all in Boston, MA.

Short title: SMOKING AND HYPERTENSION IN WOMEN

Word count= 4,496

Abstract word count=250

Dr. Bowman is supported by a Career Development Award from the Veterans Affairs  
Cooperative Studies Program. This study was supported by research grants CA-47988  
and HL-43851 from the National Institutes of Health, Bethesda, MD.

The authors report no relevant conflicts of interest pertaining to this manuscript. A full  
list of disclosures is included in the cover letter.

Corresponding author:

Thomas S. Bowman, MD MPH  
Division of Aging; Brigham and Women's Hospital  
1620 Tremont Street  
Boston, MA 02120  
Phone: 617-525-7811  
Fax: 617-525-7739  
Email: [tsbowman@partners.org](mailto:tsbowman@partners.org)

**Abstract:**

**Objectives:** To prospectively evaluate whether cigarette smoking was associated with an increased risk of developing hypertension.

**Background:** Smoking is a well-recognized risk factor for cardiovascular disease. Few prospective cohort studies have examined the relationship between smoking and hypertension.

**Methods:** We conducted a prospective cohort study among 28,236 women in the Women's Health Study who were initially free of hypertension, cardiovascular disease and cancer. Detailed risk factor information, including smoking status, was collected from self-reported questionnaires. We used Cox proportional hazards models to calculate hazard ratios (HRs) and 95% confidence intervals (CIs) of incident hypertension (defined as either new diagnosis, the initiation of anti-hypertensive medication, SBP  $\geq$ 140 or DBP  $\geq$ 90 mm Hg).

**Results:** At baseline, 51% of women were never smokers, 36% were former smokers, 5% smoked 1-14 cigarettes, and 8% smoked  $\geq$ 15 cigarettes per day. During a median of 9.8 years, there were 8,571 (30.4%) cases of incident hypertension. The age-adjusted HRs (95% CIs) of developing hypertension among never, former, and current smokers of 1-14 and  $\geq$ 15 cigarettes per day were 1.00 (reference), 1.04 (0.99-1.09), 1.00 (0.90-1.10), and 1.10 (1.01-1.19), respectively. In multivariable models further adjusting for lifestyle, clinical and dietary variables, the corresponding HRs were 1.00 (reference), 1.03 (0.98-1.08), 1.02 (0.92-1.13) and 1.11 (1.03-1.21). Among women who smoked  $\geq$ 25 cigarettes per day, the multivariable HR was 1.21 (95% CI, 1.06-1.39).

**Conclusion:** In this large cohort of women, cigarette smoking was modestly associated with an increased risk of developing hypertension, with an effect that was strongest among women smoking at least 15 cigarettes per day.

Key words: Smoking; hypertension; prospective studies; risk factors; epidemiology

### **Condensed Abstract:**

Few prospective studies have examined the relationship between smoking and hypertension. In a prospective cohort study among 28,236 women in the Women's Health Study, there were 8,571 (30.4%) cases of incident hypertension over a median follow-up of 9.8 years. The multivariable-adjusted HRs (95% CIs) of developing hypertension among never, former, and current smokers of 1-14 and  $\geq 15$  cigarettes per day were 1.00 (reference), 1.03 (0.98-1.08), 1.02 (0.92-1.13) and 1.11 (1.03-1.21). In conclusion, cigarette smoking was modestly associated with an increased risk of developing hypertension, with an effect that was strongest among women smoking at least 15 cigarettes per day.

### **Abbreviations list:**

Cardiovascular disease = CVD, Blood pressure = BP, Women's Health Study = WHS, Systolic blood pressure = SBP, diastolic blood pressure = DBP, Body mass index = BMI, Myocardial infarction = MI, Semi-quantitative food frequency questionnaire = SFFQ, Hazard ratios = HRs, Confidence intervals = CIs, World Health Organization = WHO

## **Introduction:**

Hypertension is a common, chronic condition and an improved understanding of its etiology could lead to new prevention and treatment efforts.<sup>1,2</sup> While cigarette smoking is a strong risk factor for cardiovascular disease (CVD),<sup>3</sup> its relationship with hypertension remains unclear.<sup>4</sup> Cigarette smoking causes sympathetic activation, oxidative stress and acute vasopressor effects<sup>5</sup> that are associated with increases in markers of inflammation<sup>6,7</sup> that are linked with hypertension.<sup>8</sup> Chronic smoking may also lead to endothelial dysfunction,<sup>9</sup> vascular injury, plaque progression<sup>10</sup> and increased arterial stiffness<sup>11</sup> that leads to the development of hypertension.<sup>12</sup> The long-term effect of cigarette smoking on hypertension has not been fully elucidated, and abstinence from smoking is not included in current clinical guidelines for the prevention of hypertension.<sup>13</sup>

A recent population-based study of middle-aged men reported that current cigarette smoking was independently associated with an elevated risk of developing hypertension.<sup>14</sup> However, other studies have reported lower blood pressure (BP) levels among smokers<sup>15</sup> and increases in BP after smoking cessation.<sup>16</sup> As a result, it remains unclear to what extent cigarette smoking is a risk factor for the development of hypertension.<sup>4</sup>

Therefore, we conducted a prospective cohort study among 28,236 middle-aged and older women initially free of hypertension from the Women's Health Study for the association between baseline smoking status and risk of incident hypertension.

## **Methods:**

### **Study Population**

We analyzed data from the Women's Health Study (WHS), a randomized 2x2 factorial trial of low-dose aspirin, beta-carotene and vitamin E in the primary prevention of cardiovascular disease and cancer. Between September 1992 and May 1995, letters of invitation and baseline health questionnaires were sent to more than 1.7 million female health professionals in the United States. The eligibility criteria included age  $\geq 45$  years and no history of coronary heart disease, cerebrovascular disease, cancer (except non-melanoma skin cancer) or other major illness. Additional criteria consisted of no current aspirin or non-steroidal anti-inflammatory medication use more than once a week, and no current anticoagulants or corticosteroids. A total of 453,787 women completed the questionnaires and 65,169 were willing and able to participate. These women began a three month run-in period and a total of 39,876 women who remained willing, eligible and compliant were subsequently enrolled in the trial, which had morbidity and mortality follow-up rates of 97.2% and 99.4%, respectively.<sup>17-20</sup> All participants gave written, informed consent and the study was approved by the Institutional Review Board of the Brigham and Women's Hospital in Boston, MA, USA.

Baseline information was self-reported by mailed questionnaire for demographic, lifestyle and medical history information. Every twelve months, participants were mailed follow-up questionnaires to evaluate compliance, side effects, clinical endpoints, and risk factors. For our analysis, the baseline study population was restricted to women with no self-reported pre-baseline or baseline diagnosis of hypertension, no pre-baseline or baseline antihypertensive treatment, a systolic blood

pressure (SBP) <140 mm Hg, and a diastolic blood pressure (DBP) <90 mm Hg. As a result of these exclusions, we followed 28,236 women for the development of incident hypertension.

Body mass index (BMI) at baseline was calculated as weight in kilograms divided by the square of height in meters. Participants provided information on BP from 9 ordinal SBP categories ranging from <110 mm Hg to  $\geq$ 180 mm Hg, and from 7 ordinal DBP categories ranging from <65 mm Hg to  $\geq$ 105 mm Hg. Among health professionals, self-reported BP is highly correlated with measured SBP ( $r=0.72$ ) and DBP ( $r=0.60$ ).<sup>21</sup>

#### Ascertainment of cigarette smoking status and other covariates

Cigarette smoking status was provided on the baseline questionnaire. Participants were asked to classify themselves as never smokers (<100 cigarettes in a lifetime), former smokers or current smokers. Former smokers reported whether they quit within the last year. Current smokers were categorized into 6 groups based upon number of cigarettes smoked per day (1-4, 5-14, 15-24, 25-35, 36-44 or  $\geq$ 45 cigarettes per day). Self-reported smoking status among health care professionals has been shown to be accurate in a prior meta-analysis<sup>22</sup> and in a subsequent population-based study.<sup>23,24</sup> Baseline self-reported risk factors included age, height, weight, history of high cholesterol or diabetes, exercise level, alcohol use, and parental history of myocardial infarction (MI) before age 60 years. A self-administered semi-quantitative food frequency questionnaire (SFFQ)<sup>25</sup> was also provided by participants and included information on dietary intake of fruits and vegetables<sup>26</sup> and individual energy-adjusted dietary factors such as saturated fat, sodium and potassium.<sup>27</sup>

### Evaluation of incident hypertension

Incident cases of hypertension were defined by meeting at least one of the following four criteria provided by self-reports: (1) a new physician diagnosis of hypertension on questionnaires at years 1 and 3 and then annually thereafter; (2) starting antihypertensive treatment during follow-up; (3) SBP  $\geq$ 140 mm Hg during follow-up; or (4) DBP  $\geq$ 90 mm Hg during follow-up.<sup>8</sup> Participants reporting a new hypertension diagnosis were also asked to provide the month and year that it occurred. Any missing date for incident hypertension was assigned a date by randomly selecting a date between the current and previous annual questionnaire. All participants were followed for CVD events and all relevant medical information was reviewed by the WHS Endpoints Committee composed of physicians.<sup>19</sup> Participants who developed a CVD event (including MI, stroke and revascularization) that may have affected their BP or BP management were censored at the date of diagnosis and not considered an incident case of hypertension.

### Statistical analysis

We first divided the baseline population into four groups based upon cigarette smoking status (never, past, currently smoking <15 cigarettes per day, and currently smoking  $\geq$ 15 cigarettes per day) and computed means or proportions of baseline variables for each group and for the entire cohort. Categories were compared using ANOVA or chi-square tests. We determined the number of incident hypertension cases and calculated follow-up time in person-years. To determine the effect of higher levels of

smoking, we also evaluated participants who smoked 15-24,  $\geq 25$  and  $\geq 45$  cigarettes per day.

We used Cox proportional hazards models to calculate the hazard ratios (HRs) and 95% confidence intervals (CIs) for risk of incident hypertension, using never smokers as the reference group. The crude model included only smoking status. We then adjusted for baseline age (in years). The multivariable model additionally adjusted for BMI (in  $\text{kg}/\text{m}^2$ ), exercise level (rarely/never, <1 per week, 1-3 per week,  $\geq 4$  per week), alcohol use (rarely/never, 1-3 per month, 1-6 per week,  $\geq 1$  per day), history of diabetes, history of high cholesterol  $\geq 240$  mg/dl or use of cholesterol-lowering medication, and energy-adjusted dietary intake of fruits and vegetables (servings/day), saturated fat (gm/day), sodium (mg/day) and potassium (mg/day). For exercise level and alcohol use, we did not take into account their ordinality because there is no established graded relationship between these variables and risk of hypertension. Because our variable for parental history of MI before age 60 years (yes/no) had missing data in 9.7% of participants, we included those missing this information with an indicator variable in our multivariable models. In supplementary analyses, we assessed the effect of including both SBP and DBP category in the multivariable model.

Because the relationship between smoking status and hypertension may be affected by body weight, we stratified our population by baseline BMI using World Health Organization (WHO) criteria and evaluated the risk of hypertension among normal ( $18.5 < \text{BMI} < 25 \text{ kg}/\text{m}^2$ ), overweight ( $25 \leq \text{BMI} < 30 \text{ kg}/\text{m}^2$ ) and obese ( $\text{BMI} \geq 30 \text{ kg}/\text{m}^2$ ) participants.<sup>28</sup> A test for interaction between smoking status, BMI category, and the risk

of developing hypertension was assessed where both smoking status and WHO category were ordinal variables.

We also stratified the cohort according to baseline SBP and DBP levels to determine whether smoking has differential effects on the risk of developing hypertension based upon initial BP level. The test for interaction was assessed for baseline smoking status and SBP and DBP category using ordinal variables. All HRs are presented with 95% CIs, and all reported P values are two-sided. All data analyses were performed using SAS version 9.1 (Cary, NC).

## **Results:**

Among the 28,236 women in our study, the mean ( $\pm$ SD) age at baseline was  $53.7\pm 6.6$  years, and 13.3% of participants were current smokers (**Table 1**). The mean ( $\pm$ SD) BMI was  $25.2\pm 4.5$  kg/m<sup>2</sup>, with 30% of participants classified as overweight and 13% as obese. When we stratified the cohort by smoking status, there were 14,377 never smokers, 10,115 former smokers, 1,409 participants who currently smoked 1-14 cigarettes per day, and 2,335 participants who currently smoked  $\geq 15$  cigarettes per day. Current smokers had slightly higher levels of baseline SBP ( $P < 0.01$ ) and DBP ( $P < 0.05$ ). Current smokers had a lower average BMI, higher intake of saturated fat, exercised less, and were more likely to drink alcohol (all  $P < 0.01$ ). After a median follow-up of 9.8 years, (228,933 person-years), there were 8,571 cases of incident hypertension defined by medical diagnosis (36.7%), elevations in BP (20.6%), initiation of antihypertensive treatment (3.1%), or a combination of the above (39.2%).

In Cox proportional hazard models using never smokers as the reference group, former smokers had an age-adjusted HR of 1.04 (95% CI, 0.99-1.09) and multivariable-adjusted HR of 1.03 (95% CI, 0.98-1.08) (**Table 2**). Participants who smoked 1-14 cigarettes per day had an age-adjusted HR of 1.00 (95% CI, 0.90-1.10) and multivariable-adjusted HR of 1.02 (95% CI, 0.92-1.13) while those who smoked  $\geq 15$  cigarettes per day had an age-adjusted HR of 1.10 (95% CI, 1.01-1.19) and multivariable-adjusted HR of 1.11 (95% CI, 1.03-1.21). When we considered the addition of randomized aspirin, beta-carotene and vitamin E treatment to our multivariable models, the HRs for smoking status remained unchanged. We then evaluated the 702 women who currently smoked  $\geq 25$  cigarettes per day and the 53 women who smoked  $\geq 45$  cigarettes per day, and the age-

adjusted HRs were 1.22 (95% CI, 1.07-1.39) and 1.26 (95% CI, 0.81-1.98) while the multivariable-adjusted HRs were 1.21 (95% CI, 1.06-1.39) and 1.16 (95% CI, 0.74-1.82), respectively.

We next considered the additional adjustment by baseline SBP or DBP in our multivariable models, and this only slightly affected the multivariable HRs. For example, former smokers and those currently smoking 1-14 per day had a slight increase in their HRs of hypertension, whereas the HRs among women smoking  $\geq 15$  cigarettes per day did not materially change.

When we stratified the cohort by baseline WHO category, participants with a normal BMI ( $18.5 < 25 \text{ kg/m}^2$ ) had the highest HRs of developing hypertension (P, interaction=0.04) (**Table 3**). Among these participants, the multivariable HRs for former smokers and those currently smoking 1-14 and  $\geq 15$  cigarettes per day were 1.04, 1.11 and 1.21, respectively. Among women in the normal weight category who were currently smoking  $\geq 25$  cigarettes per day, the multivariable HR was 1.31 (95% CI, 1.07-1.61). Among participants in the overweight ( $25 < 30 \text{ kg/m}^2$ ) and obese ( $\geq 30 \text{ kg/m}^2$ ) categories, there was no statistical association between cigarette smoking and hypertension.

We next stratified the cohort by baseline SBP and DBP category and evaluated smoking status and the risk of developing hypertension (**Table 4**). In age-adjusted and multivariable-adjusted models, the highest HRs were among those with a lower baseline BP. Former and current smokers with an initial DBP  $< 75$  mm Hg were at a greater relative risk of developing hypertension compared to those with a DBP that was  $\geq 75$  mm Hg (P, interaction  $< 0.01$ ). Former and current smokers with a SBP  $< 120$  mm Hg also had

higher HRs compared to those with a SBP  $\geq 120$  mm Hg, but this did not reach statistical significance (P, interaction=0.13).

## **Discussion:**

In this large prospective cohort study of 28,236 women initially free of hypertension and followed for a median of 9.8 years, we found a significantly increased risk of developing hypertension among women currently smoking at least 15 cigarettes per day. The magnitude of effect from cigarette smoking was modest, and an increased risk of hypertension was most pronounced among women with a normal BMI or low initial BP.

In the U.S., 21% of adults (44.5 million individuals) were current smokers in 2004, despite a national health objective to reduce the prevalence of smoking to less than 12% by 2010.<sup>29</sup> Among U.S. women in 2004, 18.5% were current smokers,<sup>29</sup> a proportion similar to our study cohort in which 15.3% of women were current smokers at baseline in 1992. Currently, there are at least 65 million people with hypertension in the U.S.,<sup>30</sup> and the lifetime risk of developing hypertension is estimated at 90%.<sup>31</sup> In our cohort, 30.2% of women became hypertensive over 10 years, a rate similar to that among 379 men followed for 11 years where 32.7% developed hypertension.<sup>14</sup>

While cigarette smoking has many well-established detrimental health effects,<sup>32</sup> smoking has variable effects on BP. For example, quitting smoking decreases inflammation<sup>6</sup> but may also lead to weight gain<sup>14</sup> that causes higher BPs and increased rates of hypertension.<sup>33,34</sup> Smoking may also reduce weight gain and affect the relationship between smoking and hypertension among overweight and obese women. Although the HRs of hypertension among women currently smoking 1-14 cigarettes per day did not reach statistical significance, the multivariable risk estimates were greater than 1.0 and consistent with a slightly increased risk of hypertension.

Inflammation may also mediate the relationship between smoking and high BP. In a nationally representative sample, cigarette smoking was associated with elevated inflammatory biomarkers and chronic systemic inflammation.<sup>7</sup> In the WHS, both former and current smokers had increased levels of inflammatory biomarkers,<sup>35</sup> and higher baseline levels of C-reactive protein have been associated with an increased risk of subsequent hypertension.<sup>8</sup> In a large population-based study in England, the independent association between smoking and BP was small, but causality could not be addressed in this cross-sectional study.<sup>36</sup> A prospective study among middle-aged men in Finland supported the idea that smoking may be a risk factor for hypertension, however the study size was small (n=379) and requires confirmation in other cohorts.<sup>14</sup>

There are several other potential biological mechanisms through which exposure to cigarette smoke may lead to high BP. Long-term smoking leads to abnormalities of myocardial blood flow which may be due to impairment of the vascular endothelium.<sup>37</sup> Cigarette smoking also increases BP by stiffening arteries, with particularly deleterious effects on chronic smokers.<sup>38</sup> Smoking in youth is associated with arterial changes and increased common carotid artery intima-media thickness in adulthood.<sup>39</sup> Smoking in adults is also associated with progression of atherosclerosis, and these adverse effects may be cumulative and irreversible.<sup>40</sup>

Our study has several strengths, including the prospective collection of data, large number of participants and events and long follow-up. Some potential limitations include self-reported smoking status, however both a population based study<sup>23, 24</sup> and a meta-analysis utilizing serum cotinine have confirmed that self-reports of smoking are accurate,<sup>22</sup> and self-reported medical information from health professionals is reliable.<sup>41</sup>

For self-reported hypertension in the WHS, there is an 86% validation rate which is consistent with other studies,<sup>41, 42</sup> and any potential misclassification would be random and likely bias our HRs to the null. Our participants were all apparently healthy female health professionals, possibly affecting this study's generalizability to other populations, however we have no reason to believe that the biological mechanisms by which chronic cigarette smoking may be associated with developing hypertension is unique to our study population.<sup>7, 14</sup> In the WHS, loss to follow-up was minimal regardless of smoking status and should not appreciably affect our risk estimates. We adjusted for many potential confounders in our multivariable models and while residual confounding may be present in this observational study, given the relatively small impact of adjusting for other confounders, we believe that any residual confounding would only marginally impact the HRs.

**Perspectives:**

Among apparently healthy women followed for a median of 9.8 years, an increased risk of developing hypertension was found among women currently smoking at least 15 cigarettes per day. Whether the magnitude of effect is susceptible to confounding or reflects a true association requires an improved understanding of the biological mechanisms through which smoking may lead to the development of hypertension.

Although the risk estimates were modest and found mainly among women with a normal BMI and lower initial BPs, the population-attributable risk of hypertension associated with cigarette smoking may potentially be substantial given the high prevalence of hypertension. Therefore, public health efforts to reduce the burden of smoking remain a potentially clinically relevant and effective strategy for preventing hypertension and reducing the impact of smoking on CVD.

**Acknowledgments:**

We thank the staff of the Women's Health Study for their crucial contributions, and the 39,876 dedicated and committed participants of the Women's Health Study.

## **References**

1. Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J. Global burden of hypertension: analysis of worldwide data. *Lancet* 2005;365:217-23.
2. Joint National Committee. The seventh report of the Joint National Committee on prevention, detection, evaluation, and treatment of high blood pressure. *JAMA* 2003;289:2560-72.
3. Greenland P, Knoll MD, Stamler J, et al. Major risk factors as antecedents of fatal and nonfatal coronary heart disease events. *JAMA* 2003;290:891-7.
4. Narnkiewicz K, Kjeldsen SE, Hedner T. Is smoking a causative factor of hypertension? *Blood Press* 2005;14(2):69-71.
5. Barutcu I, Esen AM, Degirmenci B, et al. Acute cigarette smoking-induced hemodynamic alterations in the common carotid artery. *Circ J* 2004;68:1127-31.
6. Bakhru A, Erlinger TP. Smoking cessation and cardiovascular disease risk factors: results from the Third National Health and Nutrition Examination Survey. *PLoS Med* 2005;2(6):e160.
7. Bazzano LA, He J, Muntner P, Vupputuri S, Whelton PK. Relationship between cigarette smoking and novel risk factors for cardiovascular disease in the United States. *Ann Intern Med* 2003;138(11):891-7.
8. Sesso HD, Buring JE, Rifai N, Blake GJ, Gaziano JM, Ridker PM. C-Reactive Protein and the risk of developing hypertension. *JAMA* 2003;290:2945-51.
9. Li H, Srinivasan SR, Chen W, Xu J, Li S, Berenson GS. Vascular abnormalities in asymptomatic, healthy young adult smokers without other major cardiovascular risk factors: the Bogalusa Heart Study. *Am J Hypertens* 2005;18:319-24.
10. Sharrett AR, Ding J, Criqui MH, et al. Smoking, diabetes, and blood cholesterol differ in their associations with subclinical atherosclerosis: the Multiethnic Study of Atherosclerosis (MESA). *Atherosclerosis* 2006;186(2):441-7.
11. Kim JW, Park CG, Hong SJ, et al. Acute and chronic effects of cigarette smoking on arterial stiffness. *Blood Press* 2005;12(2):80-5.
12. Oparil S, Zaman A, Calhoun DA. Pathogenesis of hypertension. *Ann Intern Med* 2003;139:761-76.
13. Whelton PK, He J, Appel LJ, et al. Primary prevention of hypertension. *JAMA* 2002;288:1882-8.
14. Niskanen L, Laaksonen DE, Nyyssonen K, et al. Inflammation, abdominal obesity, and smoking as predictors of hypertension. *Hypertension* 2004;44:859-65.
15. Green MS, Jucha E, Lz Y. Blood pressure in smokers and non-smokers: epidemiologic findings. *Am Heart J* 1986;111:932-40.
16. Lee D, Ha M, Kim J, Jacobs DR. Effects of smoking cessation on changes in blood pressure and incidence of hypertension. *Hypertension* 2001;37:194-8.
17. Buring JE, Hennekens CH. The Women's Health Study: summary of the study design. *J Myocardial Ischemia* 1992;4:27-9.
18. Rexrode KM, Lee IM, Cook NR, Hennekens CH, Buring JE. Baseline characteristics of participants in the Women's Health Study. *J Womens Health Gend Based Med* 2000;9:19-27.

19. Ridker PM, Cook NR, Lee I-M, et al. A randomized trial of low-dose aspirin in the primary prevention of cardiovascular disease in women. *N Eng J Med* 2005;352:1293-304.
20. Lee IM, Cook NR, Gaziano JM, et al. Vitamin E in the primary prevention of cardiovascular disease and cancer. *JAMA* 2005;294:56-65.
21. Klag MJ, He J, Mead LA, Ford DE, Pearson TA, Levine DM. Validity of physicians' self-reports of cardiovascular disease risk factors. *Ann Epidemiology* 1993;3:442-7.
22. Patrick DL, Cheadle A, Thompson DC, Diehr P, Koepsell T, Kinne S. The validity of self-reported smoking: a review and meta-analysis. *Am J Public Health* 1994;84(7):1086-93.
23. Vartiainen E, Seppala T, Lillsunde P, Puska P. Validation of self reported smoking by serum cotinine measurement in a community-based study. *J Epidemiol Community Health* 2002;56:167-70.
24. Rebagliato M. Validation of self reported smoking. *J Epidemiol Community Health* 2002;56:163-4.
25. Willett WC, Sampson L, Stampfer MJ, et al. Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol* 1985;122(1):51-65.
26. Liu S, Manson JE, Lee IM, et al. Fruit and vegetable intake and risk of cardiovascular disease: the Women's Health Study. *Am J Clin Nutr* 2000;72:922-8.
27. Willett WC. *Nutritional Epidemiology*. 2 ed. New York: Oxford University Press; 1998.
28. Leone A. Relationship between cigarette smoking and other coronary risk factors in atherosclerosis: risk of cardiovascular disease and preventive measures. *Curr Pharm Des* 2003;9(29):2417-223.
29. Centers for Disease Control and Prevention (CDC). Cigarette smoking among adults-United States, 2004. *MMWR Morb Mortal Wkly Rep* 2005;54(44):1121-4.
30. Field LE, Burt VL, Cutler JA, Hughes J, Roccella EJ, Sorlie P. The burden of adult hypertension in the United States 1999 to 2000: a rising tide. *Hypertension* 2004;44:398-404.
31. Vasani RS, Beiser A, Seshadri S, et al. Residual lifetime risk for developing hypertension in middle-aged women and men. *JAMA* 2002;287:1003-10.
32. Surgeon General. *The health consequences of smoking: a report of the Surgeon General*. Atlanta, GA: Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 2004.
33. Janzon E, Hedblad B, Berglund G, Engstrom G. Changes in blood pressure and body weight following smoking cessation in women. *J Intern Med* 2004;255:266-72.
34. Lee DH, Ha MH, Ki JR, Jacobs DR. Effects of smoking cessation on changes in blood pressure and of hypertension: a 4 year follow-up study. *Hypertension* 2001;37:194-8.
35. Bermudez EA, Rifai N, Buring JE, Manson JE, Ridker PM. Relation between markers of systemic vascular inflammation and smoking in women. *Am J Cardiol* 2002;89:1117-9.

36. Primatesta P, Falaschetti E, Gupta S, Marmot MG, Poulter NR. Association between smoking and blood pressure: evidence from the Health Survey for England. *Hypertension* 2001;37:187-93.
37. Campisi R, Czernin J, Schoder H, et al. Effects of long-term smoking on myocardial blood flow, coronary vasomotion and vasodilator capacity. *Circulation* 1998;98:119-25.
38. Kim JW, Park CG, Hong SJ, et al. Acute and chronic effects of cigarette smoking on arterial stiffness. *Blood Press* 2005;14(2):80-5.
39. Raitakari OT, Juonala M, Kahonen M, et al. Cardiovascular risk factors in childhood and carotid artery intima-media thickness in adulthood. *JAMA* 2003;290:2277-83.
40. Howard G, Wagenknecht LE, Burke GL, et al. Cigarette smoking and progression of atherosclerosis. *JAMA* 1998;279:119-24.
41. Colditz GA, Martin P, Stampfer MJ, et al. Validation of questionnaire information on risk factors and disease outcomes in a prospective cohort study of women. *Am J Epidemiol* 1986;123(5):894-9000.
42. Giles WH, Croft JB, Keenan NL, Lane MJ, Wheller FC. The validity of self-reported hypertension and correlates of hypertension awareness among blacks and whites within the stroke belt. *Am J Prev Med* 1995;11:163-9.

Table 1. Baseline characteristics of 28,236 women, according to baseline cigarette smoking status\*

Characteristic	Cigarette smoking status				Total
	Never	Past	1-14 per day	≥15 per day	
N	14,337	10,115	1,409	2,335	28,236
Age (years)†	53.6 (6.7)	54.1 (6.5)	53.6 (6.3)	53.2 (6.1)	53.7 (6.6)
BMI (kg/m <sup>2</sup> )†	25.2 (4.5)	25.3 (4.4)	24.7 (4.4)	24.9 (4.5)	25.2 (4.5)
WHO category†					
Normal	56.1	55.5	61.0	56.8	56.2
Overweight	29.3	30.5	25.3	29.1	29.5
Obese	13.4	13.0	11.9	12.0	13.1
Physical activity, %†					
Rarely/never	34.5	32.8	42.5	54.4	36.0
<1 per week	20.4	18.9	22.3	23.2	20.2
1-3 per week	33.6	34.7	28.9	19.2	32.6
≥ 4 per week	11.5	13.5	6.3	3.1	11.3
Alcohol use, %†					
Rarely/never	50.7	31.8	40.0	44.2	42.9
1-3 drinks/month	13.6	13.3	12.3	13.2	13.4
1-6 drinks/week	29.6	40.1	33.9	27.9	33.4
≥ 1 drink per day	6.1	14.8	13.8	14.8	10.3
Parental history of MI prior to age 60, %†	11.9	12.5	14.6	14.6	12.4

Diabetes, %	1.3	1.3	1.0	1.7	1.3
History of hyperlipidemia, %	24.2	25.1	23.1	26.2	24.6
Fruit and vegetables, servings/day†	6.2±3.6	6.2±3.6	5.6±3.5	4.9±3.1	6.1±3.6
Dietary saturated fat, gm/day†	19.3±4.6	19.2±4.7	20.7±4.9	22.3±5.4	19.6±4.8
Dietary sodium, gm/day†	1.86±0.33	1.87±0.34	1.85±0.35	1.90±0.38	1.87±0.34
Dietary potassium, gm/day†	3.20±0.58	3.25±0.58	3.18±0.62	3.12±0.63	3.21±0.59
SBP, mm Hg†					
<110	20.3	20.2	21.5	20.5	20.3
110-119	37.2	38.0	41.0	35.6	37.6
120-129	29.4	28.2	25.6	29.5	28.8
130-139	13.1	13.6	11.9	14.5	13.3
DBP, mm Hg‡					
<65	11.7	12.3	13.8	12.3	12.0
65-74	40.9	40.9	44.1	40.7	41.1
75-84	39.2	38.7	34.9	39.0	38.8
85-89	8.2	8.2	7.2	8.0	8.1

\* Values are mean ± standard deviation

† P<0.01

‡ P<0.05

Abbreviations: BMI=Body Mass Index; WHO=World Health Organization; SBP=Systolic Blood Pressure; DBP=Diastolic Blood Pressure

Table 2. HRs (95% CIs) from Cox proportional hazards models of incident hypertension, according to baseline cigarette smoking status among 28,236 participants

	<u>Baseline smoking status</u>				<u>Higher categories</u>	
	<u>Never</u>	<u>Former</u>	<u>1-14/day</u>	<u>≥15/day</u>	<u>15-24/day</u>	<u>≥25/day</u>
N	14,377	10,115	1,409	2,335	1,633	702
Person-years	117,020	81,898	11,523	18,491	13,120	5,371
Cases	4,273	3,150	417	731	493	238
Crude	1.00 (ref)	1.06 (1.01-1.11)	0.99 (0.90-1.10)	1.08 (1.00-1.17)	1.03 (0.94-1.13)	1.20 (1.06-1.37)
Age-adjusted	1.00 (ref)	1.04 (0.99-1.09)	1.00 (0.90-1.10)	1.10 (1.01-1.19)	1.05 (0.95-1.15)	1.22 (1.07-1.39)
Multivariable*	1.00 (ref)	1.03 (0.98-1.08)	1.02 (0.92-1.13)	1.11 (1.03-1.21)	1.07 (0.97-1.18)	1.21 (1.06-1.39)
Multivariable +SBP and DBP	1.00 (ref)	1.05 (1.00-1.10)	1.09 (0.98-1.21)	1.11 (1.02-1.20)	1.08 (0.98-1.18)	1.18 (1.03-1.35)

\* Adjusted for age, BMI, exercise level, alcohol use, parental history of MI prior to age 60 years, diabetes and high cholesterol (self-reported history or current treatment), and intake of energy-adjusted dietary factors (fruits and vegetables, saturated fat, sodium and potassium)

Note: The further consideration of randomized treatment assignment (aspirin, beta-carotene, vitamin E) in the multivariable model did not change the HRs for smoking status.

SBP=Systolic Blood Pressure; DBP=Diastolic Blood Pressure

Table 3. HRs (95% CIs) from Cox proportional hazards models\* of incident hypertension according to baseline cigarette smoking status among 28,236 participants, stratified by baseline WHO category

<u>WHO category</u>	<u>Never</u>	<u>Former</u>	<u>Current, 1-14/day</u>	<u>Current, <math>\geq</math>15/day</u>
<u>Normal (18.5-&lt;25.0 kg/m<sup>2</sup>)</u>				
N	8,058	5,616	860	1,326
Cases	1,806	1,353	219	345
Age-adjusted	1.00 (ref)	1.06 (0.99-1.14)	1.14 (0.99-1.31)	1.23 (1.09-1.38)
Multivariable*	1.00 (ref)	1.04 (0.97-1.12)	1.11 (0.96-1.28)	1.21 (1.07-1.36)
<u>Overweight (25.0-&lt;30.0 kg/m<sup>2</sup>)</u>				
N	4,213	3,088	357	679
Cases	1,522	1,149	121	235
Age-adjusted	1.00 (ref)	1.02 (0.95-1.11)	0.96 (0.80-1.16)	1.00 (0.87-1.15)
Multivariable*	1.00 (ref)	1.03 (0.95-1.12)	0.97 (0.80-1.17)	0.99 (0.86-1.14)
<u>Obese (<math>\geq</math>30 kg/m<sup>2</sup>)</u>				
N	1,927	1,313	167	281
Cases	916	627	70	141
Age-adjusted	1.00 (ref)	1.00 (0.90-1.11)	0.88 (0.69-1.12)	1.10 (0.92-1.31)
Multivariable*	1.00 (ref)	0.99 (0.89-1.11)	0.86 (0.67-1.11)	1.13 (0.94-1.36)

\* Adjusted for age, BMI, exercise level, alcohol use, parental history of MI prior to age 60 years, diabetes and high cholesterol (self-reported history or current treatment), and intake of energy-adjusted dietary factors (fruits and vegetables, saturated fat, sodium and potassium)

Table 4. HRs (95% CIs) from Cox proportional hazards models\* of incident hypertension according to baseline cigarette smoking status among 28,236 participants, stratified by baseline systolic blood pressure (SBP) and diastolic blood pressure (DBP)

	Never	Former	Current, 1-14/day	Current, ≥15/day
<u>SBP category (mm Hg)</u>				
<u>&lt;110</u>				
N	2,914	2,041	303	478
# cases	288	218	38	49
Age-adjusted	1.00 (ref)	1.06 (0.89-1.26)	1.29 (0.92-1.80)	1.03 (0.76-1.39)
Multivariable*	1.00 (ref)	1.11 (0.92-1.34)	1.33 (0.94-1.88)	1.06 (0.77-1.46)
<u>110 to 119</u>				
N	5,354	3,846	577	830
# cases	1,075	834	137	185
Age-adjusted	1.00 (ref)	1.09 (0.99-1.19)	1.21 (1.01-1.44)	1.15 (0.99-1.35)
Multivariable*	1.00 (ref)	1.05 (0.96-1.16)	1.18 (0.98-1.42)	1.15 (0.98-1.36)
<u>120 to 129</u>				
N	4,227	2,853	361	688
# cases	1,653	1,178	129	279
Age-adjusted	1.00 (ref)	1.07 (0.99-1.15)	0.89 (0.75-1.07)	1.07 (0.95-1.22)
Multivariable*	1.00 (ref)	1.09 (1.01-1.18)	0.92 (0.77-1.11)	1.09 (0.95-1.24)
<u>130 to 139</u>				
N	1,882	1,375	168	339
# cases	1,257	920	113	218
Age-adjusted	1.00 (ref)	0.99 (0.91-1.08)	1.01 (0.83-1.22)	0.99 (0.85-1.14)
Multivariable*	1.00 (ref)	0.99 (0.91-1.09)	1.05 (0.86-1.28)	1.03 (0.89-1.20)

	Never	Former	Current, 1-14/day	Current, ≥15/day
<u>DBP category (mm Hg)</u>				
<u>&lt;65</u>				
N	1,679	1,242	194	286
# cases	171	127	37	37
Age-adjusted	1.00 (ref)	0.97 (0.77-1.22)	1.98 (1.39-2.83)	1.30 (0.91-1.85)
Multivariable*	1.00 (ref)	1.06 (0.83-1.35)	2.00 (1.38-2.90)	1.23 (0.84-1.80)
<u>65 to 74</u>				
N	5,885	4,135	622	951
# cases	1,150	931	116	213
Age-adjusted	1.00 (ref)	1.16 (1.06-1.26)	0.96 (0.79-1.16)	1.22 (1.05-1.41)
Multivariable*	1.00 (ref)	1.14 (1.04-1.25)	0.96 (0.79-1.17)	1.27 (1.09-1.48)
<u>75 to 84</u>				
N	5,634	3,909	492	911
# cases	2,188	1,567	201	369
Age-adjusted	1.00 (ref)	1.03 (0.96-1.10)	1.06 (0.92-1.23)	1.08 (0.97-1.21)
Multivariable*	1.00 (ref)	1.02 (0.96-1.10)	1.10 (0.95-1.28)	1.10 (0.98-1.24)
<u>85 to 89</u>				
N	1,179	829	101	187
# cases	764	525	63	112
Age-adjusted	1.00 (ref)	0.97 (0.86-1.08)	0.92 (0.71-1.18)	0.91 (0.74-1.11)
Multivariable*	1.00 (ref)	0.97 (0.86-1.09)	0.94 (0.72-1.23)	1.01 (0.82-1.24)

\* Adjusted for age, BMI, exercise level, alcohol use, parental history of MI prior to age 60 years, diabetes and high cholesterol (self-reported history or current treatment), and intake of energy-adjusted dietary factors (fruits and vegetables, saturated fat, sodium and potassium)